

On k -cyclic SHn -algebra

A. FERNANDEZ*

Universidad de Costa Rica, 2060 San José, Costa Rica

Abstract. In this work we consider a new class of algebra called k -cyclic SHn -algebra (A, T) where A is an SHn -algebra and T is a lattice endomorphism such that $T^k(x) = x$, for all x , k is a positive integer. The main goal of this paper is to show a Priestley duality theorem for k -cyclic SHn -algebra.

Key words: De Morgan algebra, Łukasiewicz algebra, Heyting algebra, Lattices and duality.

1. Introduction

As an application to the study of switching circuits, Gr.C. Moisil introduced in [1] the symmetric Boolean algebra, that is, Boolean algebras with an automorphism of period two and later, in [2], the cyclic Boolean algebras in which the automorphism of period two is replaced by an automorphism of period k . The symmetric Boolean algebras were also studied in A. Monteiro [3, 4]. M. Abad [5] has also investigated the action of a cyclic operation on Łukasiewicz algebra, and L. Iturrioz [6] gave complete description of the variety of involutive Heyting algebra. On the other hand, it has been shown that Post and Łukasiewicz algebra are both Heyting algebra with operators. In both Łukasiewicz and Post algebra a symmetry can be expressed in terms of the primitive operations. This led to the study of more general algebra, called Symmetrical Heyting Algebras of order n (or SHn -algebra) (see [7–10]).

A k -cyclic SHn -algebra is a system (A, T) , such that A is an SHn -algebra and T is an endomorphism of A such that $T^k(x) = x$ for every $x \in A$. We usually use the same notation for a structure as for its universe. The class of k -cyclic SHn -algebra forms a variety. In this paper we give a Priestley-style duality for k -cyclic SHn -algebra.

2. Preliminaries

In this section, in order to simplify reading, we summarize the fundamental concepts we use.

Recall that V. Sofronie-Stokkermas [11] introduce the category **SHn** of SHn -algebra and SHn -homomorphisms, where an SHn -algebra is an algebra $\langle A, \wedge, \vee, \Rightarrow, \sim, 0, 1, S_1, \dots, S_{n-1} \rangle$ such that $\langle A, \wedge, \vee, \Rightarrow, \sim, 0, 1 \rangle$ is a symmetric Heyting algebra (see [12]) and S_1, \dots, S_{n-1} are unary operators defined on A fulfilling the following equalities:

- (S1) $S_i(a \wedge b) = S_i a \wedge S_i b$,
- (S2) $S_i(a \Rightarrow b) = (\bigwedge_{k=i}^n S_k(a) \Rightarrow S_k(b))$,
- (S3) $S_i(S_j(a)) = S_j(a)$, for every $i, j = 1, \dots, n - 1$,
- (S4) $S_1(a) \vee a = a$,

- (S5) $S_i(\sim a) = \sim S_{n-i}(a)$, for $i = 1, \dots, n - 1$,
- (S6) $S_1(a) \vee \neg S_1(a) = 1$, with $\neg a = a \Rightarrow 0$.

In addition, this author extended Esakia duality [13, 14] to the category **SHnSp** whose objects are SHn -spaces and whose morphisms are SHn -functions. Specifically, an SHn -space is a system $(X, \leq, \tau, s_1, \dots, s_{n-1}, g)$ such that (X, \leq, τ) is an Esakia space and for every $x, y \in X$ the following conditions are satisfied:

- (1) if $x \leq y$ then $g(y) \leq g(x)$,
- (2) $g(s_i(x)) = s_{n-i}(g(x))$,
- (3) $g(g(x)) = x$,
- (4) $s_j(s_i(x)) = s_j(x)$,
- (5) $s_1(x) \leq x$,
- (6) $x \leq s_{n-1}(x)$,
- (7) $s_i(x) \leq s_j(x)$ for every $i \leq j$,
- (8) if $x \leq y$ then $s_i(x) = s_i(y)$,
- (9) for all $i = 1, \dots, n - 1$, $x \leq s_i(x)$ or $s_{i+1}(x) \leq x$.

and an SHn -function from an SHn -space $(X, \leq, \tau, s_1, \dots, s_{n-1}, g)$ into another $(X', \leq', \tau', s'_1, \dots, s'_{n-1}, g')$ is an Esakia morphism (continuous bounded morphism) $f : X \rightarrow X'$ such that preserve the operations s_1, \dots, s_{n-1}, g . Besides, he proved that **SHn** is dually equivalent to **SHnSp**.

3. Priestley duality for k -cyclic SHn -algebra

In this section we study SHn -algebra endowed with a lattice endomorphism, more precisely:

Definition 1. An k -cyclic SHn -algebra is a pair (A, T) where A is an SHn -algebra and T is a lattice endomorphism which satisfies the following properties:

- (K) $T^k(x) = x$, for every $x \in A$.

By **kSHn** we will denote the category of k -cyclic SHn -algebra and their corresponding homomorphisms.

We indicate a Priestley-style duality for k -cyclic SHn -algebra.

*e-mail: arielfernandez@gmail.com

Definition 2. A k -space is a pair (X, t) where X is an SHn -space and the following conditions are satisfied:

- (k1) $t : X \rightarrow X$ is an homeomorphism,
- (k2) $t : X \rightarrow X$ is an order-isomorphism,
- (k3) $t^k(x) = x$, for every $x \in X$.

Let \mathbf{kSp} be the category whose objects are k -spaces and whose morphisms are SHn -functions such that preserve the operation t .

For every k -cyclic space (X, t) , let $E(X)$ be the set of clopen order-filters of X . On $E(X)$ the following operations can be defined:

- (a) $\vee : E(X) \times E(X) \rightarrow E(X)$ is defined by $U \vee V := U \cup V$.
- (b) $\wedge : E(X) \times E(X) \rightarrow E(X)$ is defined by $U \wedge V := U \cap V$.
- (c) $\Rightarrow : E(X) \times E(X) \rightarrow E(X)$ is defined by $U \Rightarrow V := \{x \in X \mid x \leq y \text{ and } y \in U \text{ implies } y \in V\}$.
- (d) $\sim : E(X) \times E(X) \rightarrow E(X)$ is defined for every $U \in E(X)$ by $\sim U := X \setminus g^{-1}(U)$.
- (e) For every $i = 1, \dots, n-1$, $S_i : E(X) \rightarrow E(X)$ is defined for every $U \in E(X)$ by $S_i(U) := s^{-1}(U)$.
- (f) $T : E(X) \rightarrow E(X)$ is defined for every $U \in E(X)$ by $T(U) := t^{-1}(U)$.

For every morphism of k -cyclic space $\varphi : X \rightarrow X'$ let $E(\varphi) : E(X') \rightarrow E(X)$ be defined by $E(\varphi)(U) = \varphi^{-1}(U)$ for every $U \in E(X')$.

Proposition 1. The functor $E : \mathbf{kSp} \rightarrow \mathbf{kSHn}$ is well-defined, i.e. the following holds:

- (i) The algebra $(E(X), T)$ is an k -cyclic SHn -algebra.
- (ii) If $\varphi : X \rightarrow X'$ is a morphism of k -spaces, then $E(\varphi) : E(X') \rightarrow E(X)$ is a morphism of k -cyclic SHn -algebra.

Proof. We shall only prove that for all $U \in E(X)$, (t4) is satisfied. Indeed, the following conditions are equivalents:

- (1) $x \in T^k(U)$,
- (2) $t^k(x) \in U$,
- (3) $x \in U$.

For every k -cyclic SHn -algebra (A, T) let $D(A) = (PF(A), \leq, \tau, s_1, \dots, s_{n-1}, g, t)$ where $(PF(A), \leq, \tau, s_1, \dots, s_{n-1}, g)$ is the Priestley space of the SHn -algebra A and $t : D(A) \rightarrow D(A)$ is defined for every $F \in D(A)$ by $t(F) = T^{-1}(F)$.

For every morphism of k -cyclic SHn -algebra $f : A \rightarrow A'$ let $D(f) : D(A') \rightarrow D(A)$ be defined by $D(f)(F) = f^{-1}(F)$ for every $F \in D(A')$.

Proposition 2. The functor $D : \mathbf{kSHn} \rightarrow \mathbf{kSp}$ is well-defined, i.e. the following holds:

- (i) For every k -cyclic SHn -algebra (A, T) , $(D(A), t)$ is an k -space.

- (ii) If $f : A \rightarrow A'$ is a morphism of SHn -algebra, then $D(f) : D(A') \rightarrow D(A)$ is a morphism of k -spaces.

Proof. We can only prove that for all $F \in D(A)$, (K) is satisfied. Indeed, the following conditions are equivalents:

- (1) $x \in t^k(F)$,
- (2) $T^k(x) \in F$,
- (3) $x \in F$.

From Proposition 1 and 2 and taking into account the results indicad in [11] we have.

Theorem 1. The categories \mathbf{kSHn} and \mathbf{kSp} are dually equivalent.

REFERENCES

- [1] Gr. C. Moisil, "Algebra schemelor cu elemente ventil", *Revista Universitatii C. I. Parhon si a Politehnicii Bucuresti Seria St. Nat.* 4-5, 9-42 (1954).
- [2] Gr. C. Moisil, *Essais sur les Logiques Non Chrysippiennes*, Academiei Bucurest, Bucurest, 1972.
- [3] A. Monteiro, *Algebras de Boole Involutivas*, Universidad Nacional del Sur, Bahía Blanca, 1969.
- [4] A. Monteiro, *Algèbres de Boole Cycliques*, *Revue Roumaine de Math. Pures et Appliquées* 23, 71-76 (1978).
- [5] M. Abad, "Estructuras cíclica y monádica de un álgebra de Łukasiewicz n -valente", *Notas de Lógica Matemática* 36, CD-ROM (1988).
- [6] L. Iturrioz, "Algèbres de Heyting trivalentes involutives", *Notas de Lógica Matemática* 31, CD-ROM (1974).
- [7] L. Iturrioz, "Łukasiewicz and symmetrical Heyting algebra", *Zeitschrift für Mathematische Logik und Grundlagen der Mathematik* 23, 131-136 (1977).
- [8] L. Iturrioz, *Modal Operators on Symmetrical Heyting Algebra*, Universal Algebra and Applications, Banach Center Publications 9, ed. T. Traczyk, pp. 289-303, PWN-Polish Scientific Publishers, Warsaw, 1982.
- [9] L. Iturrioz, "Symmetrical Heyting algebra with operators", *Zeitschrift für Mathematische Logik und Grundlagen der Mathematik* 29, 33-70 (1983).
- [10] L. Iturrioz and V. Sofronie-Stokkermans, " SHn -algebra (abbreviation of symmetrical Heyting algebra of order n)", in: *Atlas of Many-Valued Structures, COST Action 15*, eds. L. Iturrioz, E. Orłowska, E. Turunen, pp. 1235-9599, Tampere Univ. of Technology, Tampere, 2000.
- [11] V. Sofronie-Stokkermans, "Priestley duality for SHn -algebras and applications to the study of Kripke-style models for SHn -logics", *Multiple Valued Logic, Int. J.* 5 (4), 281-305 (1999).
- [12] A. Monteiro, "Sur les algèbres de Heyting symétriques", *Portugaliae Mathematica* 39, 1-237 (1980).
- [13] L.L. Esakia, "Topological Kripke models", *Soviet Math. Dokl.* 15, 147-151 (1974).
- [14] L.L. Esakia, "The problem of dualism in the intuitionistic logic and Brouwerian lattices", *V Int. Congress Logic, Methodology and Philosophy of Science* 1, 7-8 (1975).